Powerful Outflows and Feedback from Active Galactic Nuclei

Andrew King and Ken Pounds



Presented by Lukas Zalesky

Understanding the Influence of SMBH

Three Important Equations

1. $M_{SMBH} \sim 10^{-3} \times M_{Bulge}$

2. $M_{SMBH} \approx 3 \times 10^{8} M_{\odot} \sigma_{200}^{\alpha}$

3. $R_{inf} \approx 8\ M_8/\sigma_{200}^{2}$ pc

How can this be so?

Significance

1. The mass of the SMBH is almost always a constant fraction of the mass of the bulge


1. The mass of the SMBH is tightly correlated to the velocity dispersion in the bulge

2. Most of the entire galaxy is gravitationally unaware of the SMBH

Methods of Communication

Radiation: how all the accretion energy is initially released

Mechanical, i.e., literally moving gas around: huge luminosities drive gas through the galaxy




IRAS F11119+3257 artist depiction

<u>Jets vs. Winds</u>


Jets: highly collimated flows driven from very near the SMBH, somewhat rare

Winds: nearly isotropic outflows, common to many AGN

Detecting Winds: Spectrum of PG1211+143



Modeling Winds: Spectrum of PG1211+143




Gaussian fit, v ~ 0.12c

Photoionized gas model, v ~ 0.15c

Observational Evidence of UFOs

(Ultra-Fast Outflows)





Cappi 2006 found 7 non-BAL objects with v ~ 0.1c

Tombesi et al. 2010 found outflows in 15/42 radio-quiet objects, v ~ [0.1c,0.3c] (XMM-Newton archive)

Gofford et al. 2013 found similar outflows in 20/51 AGN, also with v ~ [0.1c,0.3c] (Suzaku archive)

→ Rather common

Energy Output and Binding Energies

$$E_{\rm BH} \simeq \eta M c^2 \sim 2 \times 10^{61} M_8 \ {\rm erg}$$

$$E_{\rm bulge} \sim M_{\rm b} \sigma^2 \sim 8 \times 10^{58} M_8 \sigma_{200}^2 \ {\rm erg}$$

Black hole mass should be related to galaxy properties that depend on galactic gas, since it is through accretion of this gas that it grows

The important comparison is between E_{BH} and E_{gas}


$$E_{\rm gas} = f_{\rm g} E_{\rm bulge},$$

$$E_{\rm BH} \sim 2 \times 10^3 E_{\rm gas}$$

The energy is more than enough to disrupt all of the gas in the bulge of galaxy, so it must not be communicated efficiently → slowed outflows

Shocked Outflows

$$E_{\rm mom} \sim \frac{\sigma}{c} E_{\rm BH\ wind} \sim \frac{\sigma}{c}\frac{\eta^2}{2} Mc^2 \sim 5 \times 10^{-5} Mc^2 \sim 0.1 E_{\rm gas}$$



Significant Inverse Compton cooling


Energy Dominated Flows

•Cooling time is not significant

$$E_{\rm wind} \simeq 0.05 L_{\rm Edd} \simeq 100 E_{\rm gas}$$

•Expansion is adiabatic so it retains its heat, and at greater distances the cooling time begins to increase, confining momentum flows to smaller radii and allowing energy dominated flows to extend much further

•Could not be supported long term because this would blow out all of the gas that the SMBH feeds on → couldn't reach observed masses


Shock front
(momentum and
pressure conserved)
Wind shock
(thermalized energy rapidly
lost to cooling, only ram
pressure conserved)
Fast wind
($\dot{E}_w = \eta L_{Edd}/2$)
ISM
Momentum-driven outflow
ISM
Cooling ISM
Cold shocked ISM
($\dot{E} \approx \alpha L_{Edd}/c \ll \dot{E}_w$)
Cold shocked wind
($\dot{E} \approx \alpha L_{Edd}/c \ll \dot{E}_w$)
Cooling wind
Fast wind
ISM shock
(thermalized energy
rapidly lost to cooling,
only ram pressure conserved)
SMBH
Contact discontinuity
(momentum and
pressure conserved)
Energy-driven outflow
Fast wind
($\dot{E}_w = \eta L_{Edd}/2$)
Shocked,
adiabatically expanding
wind
($\dot{E}_w = \eta L_{Edd}/6$)
Shocked,
adiabatically expanding
ISM
($\dot{E}_w = \eta L_{Edd}/3$)
ISM
Wind shock
(energy conserved)
ISM shock
(energy conserved)

The M-σ Relation

A quick derivation that leads to a qualitatively similar M-σ relation

$$\rho(r) = \frac{f_g \sigma^2}{2\pi G r^2},$$

➔

$$M(R) = \frac{2\sigma^2 R}{G}.$$

$$M_g(R) = \frac{2 f_g \sigma^2 R}{G}$$

➔

$$\frac{d}{dt}[M_g(R)\dot{R}] + \frac{G M_g(R)[M + M(R)]}{R^2} = \frac{L_{Edd}}{c},$$

$$\frac{d}{dt}(R\dot{R}) + \frac{GM}{R} = -2\sigma^2\left(1 - \frac{M}{M_\sigma}\right),$$

where

$$M_\sigma = \frac{f_g \kappa}{\pi G^2}\sigma^4.$$

$$R^2\dot{R}^2 = -2GMR - 2\sigma^2\left[1 - \frac{M}{M_\sigma}\right]R^2 + \text{constant}.$$

➔

$$\dot{R}^2 \simeq -2\sigma^2\left[1 - \frac{M}{M_\sigma}\right],$$

What if $M < M_\sigma$?

What happens at M_σ and Beyond

$M_{SMBH} \approx 3 \times 10^8 M_\odot \sigma_{200}^{\alpha}$

$$M_\sigma = \frac{f_g \kappa}{\pi G^2} \sigma^4 \simeq 3.2 \times 10^8 \, M_\odot \sigma_{200}^4$$

Something must happen to halt black hole growth at this point → The black hole must expel all the gas

For $M \gg M_\sigma$, the solution to ----------> allows for R to grow very large, larger than the critical radius at which cooling takes place.

$$\dot{R}^2 \simeq -2\sigma^2 \left[1 - \frac{M}{M_\sigma}\right],$$

For $M \gg M_\sigma$, the shock wave becomes energy driven, and can extend for kpc scales → Clear out

The Effects of a "Clear out" Considering Galactic Morphology

Disk Galaxies, e.g. spirals

Eventually the outflow must encounter the galactic disk. The resistance causes the flow proceed above and below the disk, however, $P_{outflow} >> P_{disk}$


→ Triggered star formation

Elliptical Galaxies

In the absence of a disk, the energetic outflows proceed more or less uninterrupted, leaving spherical galaxies "red and dead"

Different morphologies and locations in which galaxies are found will yield different interactions between galaxies and their environments →
Three (four) M-σ relationships

Three (Four) M-σ Relationships



Black Hole Mass – Bulge Mass Relation

Our previous discussion asserts that σ determines M_{SMBH}, so we cannot say that M_{bulge} independently determines M_{SMBH}.

Additionally, M_{SMBH} cannot determine M_{bulge} because it is in the form of stars

Hypothesis: Whatever determines M_{bulge} also makes it proportional to σ^4 (and therefore M_{SMBH}). Why?

Star Formation & Stellar Feedback

$$L_* \sim 2 \times 10^{10}\, \mathrm{L}_\odot\, \sigma_{200}^4.$$

M/L ~ 5 (Faber-Jackson is empirical)

$$M_* \sim 1 \times 10^{11}\, \mathrm{M}_\odot \sigma_{200}^4 \sim 10^3 M.$$

$$M_\mathrm{b} \sim \frac{0.14 f_\mathrm{g} t_\mathrm{H} \sigma^4}{\epsilon_* c\, G},$$

$$M \simeq M_\sigma \sim \frac{1.8 \kappa \epsilon_* c}{\pi G t_\mathrm{H}} M_\mathrm{b} \sim 10^{-3} M_\mathrm{b},$$

A Statistical Interpretation

The Less Accepted Hypothesis:

Jahnke & Maccio (2011) argue that the observed scaling relationships are the consequence of repeated mergers

M_{SMBH} is proportional to M_{bulge} simply due to the central limit theorem, i.e., a value that centers more and more closely on the mean of the system for increasing numbers of mergers.

Counter-Argument:

1. This hypothesis does not determine the ratio of M_{SMBH}/M_{bulge}

2. This doesn't simultaneously explain both M_{SMBH}-M_{bulge} and M_{SMBH}-σ

3. It would be incredible coincidence that this "chance" collision process produces the same M_{SMBH}-σ relation needed to balance the weight of the bulge gas

Summarizing Wind-Momentum Driving Effects ~ Small Scale

1. Super-solar elemental abundances in observed AGN spectra

2. Dark matter cusp removal

3. Quiescence of AGN hosts

Summarizing Wind-Energy Driving Effects ~ Large Scale

1. Metals in the circumgalactic medium

2. Mechanical luminosities of galaxy-scale molecular outflows

3. Suppression of cosmological infall

Closing Remarks

Remaining Question at Hand:

What mechanism can produce such a ready supply of gas with very small angular momentum, such that it can accrete onto the central SMBH fast enough for it to grow relatively quickly?

→ Not gravity

→ Perhaps feedback from SMBH can cause this?